


# Guide to Internal Parasites of Ruminants

Sponsored by Intervet—providers of Panacur®/Safe-Guard® to the livestock industry.

This guide intentionally obscured  
You will get an unobscured copy  
with purchase.



**Trichostrongylus axei**

**Cyathostomum**

**Haemonchus contortus**

**Moneilema**

**Haemonchus contortus**



**Haemonchus contortus**

**Haemonchus contortus**

**Trichostrongylus axei**

**Cooperia oncophora**



**Trichostrongylus axei**

**Strongylidae**

**Coccidia**

**Dictyocaulus viviparus**

**Microsporidia**

# Health Impact and Characteristics of Internal Parasites

Parasite	Approximate Length (µm)	Characteristics
<b>Ostertagia</b> (brown stomach worm) ▪ Suppresses appetite ▪ Weight loss, poor body condition	60-70	Medium-sized, standard strongyle egg; barrel-shaped sidewalls; large number of blastomeres nearly fills egg
<b>Haemonchus</b> (barberpole worm) ▪ Anemia ▪ Death common in sheep and goats	85	Larger and rounder than <i>Ostertagia</i> egg; blastomeres more easily seen than in <i>Ostertagia</i>
<b>Trichostrongylus</b> (bankrupt worm) ▪ Watery diarrhea ▪ Slowed growth	85	Often shaped like a kidney bean; one side is more rounded than the other; there is usually a lot of clear space within the egg
<b>Cooperia</b> (small intestinal worm) ▪ Diarrhea ▪ Slowed growth	75-85	Medium-sized egg with parallel sides and numerous blastomeres that are hard to distinguish
<b>Nematodirus</b> (threadneck worm) ▪ Diarrhea ▪ Slowed growth ▪ Emaciation and death in sheep, young cattle	200	Large egg; looks like an American football with basketballs inside; two to eight large blastomeres are surrounded by a fluid-filled cavity
<b>Oesophagostomum</b> (nodular worm) ▪ Diarrhea ▪ Slowed growth	95	Medium-sized to large egg; about one and a half times the size of the <i>Ostertagia</i> egg; 16 to 32 blastomeres; are easier to see than those of <i>Haemonchus</i>
<b>Bunostomum</b> (hookworm) ▪ Anemia ▪ Weight loss	100	Medium-sized to large egg; four to eight blastomeres; sometimes the walls are thick and rectangular
<b>Strongyloides</b> (threadworm) ▪ Diarrhea in young ▪ Fatal infections reported in young raised on sawdust	40-65	Small egg with a thin shell containing an L1 larva that can be seen under low power
<b>Trichuris</b> (whipworm) ▪ Reduced appetite ▪ Slowed growth	75	Egg is shaped like an American football and has two protruding polar caps; the shell is double and thick
<b>Coccidia</b> ▪ Bloody diarrhea in young cattle ▪ Reduced weight gains in sheep, goats, cattle ▪ Death can occur in calves, lambs, kids and adult goats	16-47	Coccidia appear small in size, pink in color; size and shapes vary depending on species
<b>Moniezia</b> (tapeworm) ▪ Blocks small intestine in lambs ▪ Loss of nutrients	80x80	Quadrangular, somewhat irregular; contains a circular or pear-shaped apparatus at one end
<b>Dictyocaulus</b> (lungworm) ▪ Cough ▪ Reduced appetite and milk production	450	Rectal sample of feces needed for positive identification; L1 larva found in feces; flattened head and tail end in blunt point

# Modified Wisconsin Sugar Fecal Worm Egg Flotation Method

1. Fecal samples can be stored for long periods if refrigerated (not frozen).
2. Slides can usually be placed in the refrigerator for several days prior to reading.
3. Identify parasites present:  
+ (1-10 eggs/sample) ++ (11-50 eggs/sample) +++ (over 50 eggs/sample)
4. # of eggs found x 150 = # of eggs per pound feces

### 5. Materials needed:

- a. Sugar solution plus dispensing bottle, gun, or syringe
- b. Tea strainer
- c. 3 oz. and 5 oz. Dixie cups
- d. Tongue depressors
- e. Taper-bottom test tubes
- f. Test tube rack
- g. Standard microscope slides
- h. Centrifuge
- k. Microscope